

Modelling dependence in finance using copulas*

Statistics 2001, Concordia University, Montréal†

Thierry Roncalli

Groupe de Recherche Opérationnelle
Crédit Lyonnais

*These slides may be downloaded from <http://gro.creditlyonnais.fr> (the direct link is <http://gro.creditlyonnais.fr/content/wp/copula-stat2001-canada.pdf>).

†I would like to thank Professor Christian Genest for his invitation.

Agenda

1. The Gaussian assumption in finance
2. Copulas and multivariate financial models
3. An open field for risk management
 - Market risk
 - Operational risk
 - Credit risk
4. New pricing methods with copulas
 - Multi-asset options
 - Credit derivatives

1 The Gaussian assumption in finance

We consider the ‘universal’ financial model. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space. The asset prices processes $S_1(t)$ and $S_2(t)$ are given by the SDE representation

$$\begin{cases} dS_1(t) = \mu_1 S_1(t) dt + \sigma_1 S_1(t) dW_1(t) \\ dS_2(t) = \mu_2 S_2(t) dt + \sigma_2 S_2(t) dW_2(t) \end{cases}$$

where $W_1(t)$ and $W_2(t)$ are two \mathcal{F}_t -brownian motions with

$$\mathbb{E} [W_1(t) W_2(t) \mid \mathcal{F}_{t_0}] = \rho(t - t_0)$$

It comes that the logarithm return of assets is gaussian (= Gaussian assumption in finance).

Empirical facts: **Asset returns are not gaussian** (see the financial econometric literature on ARCH, long-memory, Lévy processes, etc.).

Problem: Univariate financial models are not gaussian, but multivariate financial models are gaussian!

2 Copulas and multivariate financial models

1. How to define multivariate financial models compatible with univariate non-Gaussian financial models?

Copula = a powerful tool

2. How to obtain tractable multivariate financial models (in terms of computational time)?
3. How to specify multivariate financial models which may be understood/used by the finance industry?

Copula = a promising tool

Copulas have been already incorporated in some software solutions:

- SAS Risk Dimensions
- Palisade @Risk

2.1 Pearson correlation and dependence

Pearson correlation ρ = linear dependence measure.

For two given asset prices processes $S_1(t)$ and $S_2(t)$ which are GBM, the range of the correlation is

$$\rho^- \leq \rho(S_1(t), S_2(t)) \leq \rho^+$$


with

$$\rho^\pm = \frac{\exp(\pm \sigma_1 \sigma_2 (t - t_0)) - 1}{\sqrt{\exp(\sigma_1^2 (t - t_0)) - 1} \cdot \sqrt{\exp(\sigma_2^2 (t - t_0)) - 1}}$$


$\rho(S_1(t), S_2(t)) = \rho^-$ (resp. ρ^+) $\Leftrightarrow C\langle S_1(t), S_2(t) \rangle = C^-$ (resp. C^+) $\Leftrightarrow S_2(t) = f(S_1(t))$ with f a decreasing (resp. increasing) function

$\text{Perfect dependence} \neq |\rho| = 1$


$\sigma_1 = 0.1 \quad (t-t_0) = 1$




$\sigma_1 = 1 \quad (t-t_0) = 1$



$\sigma_1 = 2 \quad (t-t_0) = 1$



$\sigma_1 = 0.75 \quad \sigma_2 = 1.25$



Permissible range of $\rho(S_1(t), S_2(t))$
when $S_1(t)$ and $S_2(t)$ are two GBM processes

2.2 Copula: a new tool in finance

- introduced by Embrechts et al. [14].
- Market risk: Bouyé et al. [2], Cherubini et al. [8], Durrleman et al. [13], Embrechts et al. [15], Luciano et al. [24], Tibiletti [33].
- Credit risk: Coutant et al. [11], Frey et al. [17], Georges et al. [18], Giesecke [19], Hamilton et al. [20], Lindskog et al. [23], Maccarinelli et al. [25].
- Operational risk: Ceske et al. [5] [6], Frachot et al. [16].
- Asset prices modelling: Bouyé et al. [4], Malavergne et al. [26], Patton [27], Rockinger et al. [28], Scaillet [31].
- Credit derivatives pricing: Li [21], Georges et al. [18], Schönbucher et al. [32].
- Multi-asset options pricing: Bikos [1], Cherubini et al. [7], Coutant et al. [10], Durrleman [12], Rosenberg [29] [30].

2.3 Copulas in a nutshell

A copula function \mathbf{C} is a multivariate probability distribution with uniform $[0, 1]$ margins.

$\mathbf{C}(\mathbf{F}_1(x_1), \dots, \mathbf{F}_N(x_N))$ defines a multivariate cdf \mathbf{F} with margins $\mathbf{F}_1, \dots, \mathbf{F}_N \Rightarrow \mathbf{F}$ is a probability distribution with given marginals.


The copula function of the random variables (X_1, \dots, X_N) is **invariant** under strictly increasing transformations ($\partial_x h_n(x) > 0$):

$$\mathbf{C}\langle X_1, \dots, X_N \rangle = \mathbf{C}\langle h_1(X_1), \dots, h_N(X_N) \rangle$$


... the copula is invariant while the margins may be changed at will, it follows that is precisely the copula which captures those properties of the joint distribution which are invariant under a.s. strictly increasing transformations (Schweizer and Wolff [1981]).

\Rightarrow **Copula = dependence function of r.v.** (Deheuvels [1978]).


$F_1 = \text{IG}(2, 1.5)$




$F_2 = \text{Beta}(2, 2)$



PDF of the Copula



PDF of $F(x_1, x_2) = C(F_1(x_1), F_2(x_2))$



Bivariate distribution with given marginals

2.4 The Normal copula

Two characteristics in finance: High dimensional problems (e.g. a portfolio with 1000 securities) and probabilistic properties of the models (e.g. markovian property).

All the copula functions are not good candidates for financial application in an industry point of view.

The Normal copula has not yet been extensively studied (see however Song [2000]). Nevertheless, it may be an 'industrial' copula.

Remark 1 *The multivariate normal distribution is very tractable. It is very easy to estimate the parameters and simulation is straightforward. Moreover, this distribution has nice properties.*

Is it also the case for the Normal copula?

The copula function

$$C(\mathbf{u}; \rho) = \Phi\left(\Phi^{-1}(u_1), \dots, \Phi^{-1}(u_N); \rho\right)$$

The density is

$$c(\mathbf{u}; \rho) = |\rho|^{-\frac{1}{2}} \exp\left(-\frac{1}{2} \varsigma^\top (\rho^{-1} - \mathbb{I}) \varsigma\right)$$

with $\varsigma = (\Phi^{-1}(u_1), \dots, \Phi^{-1}(u_N))$.

The Ψ transform We define the operator Ψ as follows

$$\begin{aligned} \Psi[\mathbf{F}] &: \mathbb{R} \longrightarrow \mathbb{R} \\ x &\longmapsto \Psi[\mathbf{F}](x) = \Phi^{-1}(\mathbf{F}(x)) \end{aligned}$$

We note also Ψ^{-1} the (left) inverse operator ($\Psi^{-1} \circ \Psi = 1$), i.e. $\Psi^{-1}[\mathbf{F}](x) = \mathbf{F}^{[-1]}(\Phi(x))$.

Estimation The log-likelihood function is

$$\ell(\mathbf{u}; \rho) = -\frac{T}{2} \ln |\rho| - \frac{1}{2} \sum_{t=1}^T \varsigma_t^\top (\rho^{-1} - \mathbb{I}) \varsigma_t$$

and the ML estimate of ρ is also $\hat{\rho}_{\text{ML}} = \frac{1}{T} \sum_{t=1}^T \varsigma_t^\top \varsigma_t$.

Two-stage (Joe and Xu [1996]) and omnibus (Genest, Ghoudi and Rivest [1995]) estimators can then be obtained with

$\varsigma_t = (\psi[\mathbf{F}_1](x_1^t), \dots, \psi[\mathbf{F}_N](x_N^t))$:

1. IFM estimate: $\mathbf{F}_n = \text{MLE of the } n^{\text{th}} \text{ marginal distribution.}$
2. Omnibus estimate : $\mathbf{F}_n = n^{\text{th}} \text{ empirical distribution.}$

\Rightarrow The data are mapped to uniforms and transformed with the inverse gaussian distribution. The correlation parameter ρ of the Normal copula is then equal to the Pearson product moment of the transformed data.

Simulation

- Generate a gaussian vector \mathbf{v} of random variables with correlation ρ .
- To simulate a vector \mathbf{x} of random variables with marginals $\mathbf{F}_1, \dots, \mathbf{F}_N$ and a Normal copula with parameters ρ , we use the following transformation

$$\mathbf{x} = \left(\psi^{-1} [\mathbf{F}_1] (v_1), \dots, \psi^{-1} [\mathbf{F}_N] (v_N) \right)$$

Application to marketing Segmentation is a useful tool for marketing (and scoring). Let Y be a random variable which corresponds to the target. The main idea is to define classes

| Class | Definition |
|----------|---------------------------|
| 1 | $Y \leq y_1$ |
| \vdots | |
| M | $y_{M-1} \leq Y \leq y_M$ |

For example, let Y be a (potential) rentability index. The bank would like to capture the most profitable customers. It can define the targets y_m by a quantile rule $\Pr\{Y \leq y_m\} = \tau_m$. For each class, it will define a specific customer relationship policy. For example, it will decide how much to spend on capture customers for a specific class.

One statistical tool which are used is the linear quantile regression

$$Y_n = X_n^\top \beta + u_n$$

where X_n are the characteristics of the individual n . If we reformulate the linear regression with only positive terms:

$$Y_n = X_n^\top \beta + u_n = \sum_{k=1}^K x_{n,k} (\beta_n^+ - \beta_n^-) + u_n^+ - u_n^-$$

We can show that the solution of the quantile regression $\Pr \{Y \leq y_m\} = \tau_m$ may be found using linear programming:

$$z = \arg \min c^\top z$$

$$\text{u.c.} \quad \begin{cases} Az = y \\ z \geq 0 \end{cases}$$

where $X = (X_1, \dots, X_N)^\top$, $A = (X, -X, I_N, -I_N)$, $y = (Y_1, \dots, Y_N)^\top$, $z = (\beta^+, \beta^-, u^+, u^-)^\top \in \mathbb{R}^{2K+2N}$ and $c = (\mathbf{0}, \mathbf{0}, \tau_m \mathbf{1}, (1 - \tau_m) \mathbf{1})^\top$.

Computational issues ($\dim A \simeq N \times 2N$) = very large-scale problem. Portnoy and Koenker [1997] suggest then to use an interior-point method.

Problem: how to proceed when (Y, X) are not gaussian? One solution is to assume that only the copula of (Y, X) is Normal. In this case, we can use the Portnoy-Koenker algorithm with the transformed variables $Y_i = \Psi[\mathbf{F}_i](X_i)$.

Let consider the bivariate case*. We have $\partial_1 C(u_1, u_2) = \Phi(\varsigma)$ where $\varsigma = (1 - \rho^2)^{-\frac{1}{2}} [\Phi^{-1}(u_2) - \rho\Phi^{-1}(u_1)]$. The relationship between u_2 and u_1 in the expression $\Pr\{U_2 \leq u_2 \mid U_1 = u_1\} = \tau$ is also given by

$$u_2 = \Phi\left(\rho\Phi^{-1}(u_1) + \sqrt{1 - \rho^2}\Phi^{-1}(\tau)\right)$$

If the margins are gaussian, we obtain the well-known curve


$$X_2 = \left[\mu_2 - \rho\frac{\sigma_2}{\sigma_1}\mu_1 + \sqrt{1 - \rho^2}\Phi^{-1}(\tau)\right] + \rho\frac{\sigma_2}{\sigma_1}X_1$$

We remark that the relationship is **linear**. When the margins are not gaussian, the relationship is linear in the Ψ projection space:


$$\Psi[\mathbf{F}_2](X_2) = a + b\Psi[\mathbf{F}_1](X_1)$$

where $a = \sqrt{1 - \rho^2}\Phi^{-1}(\tau)$ and $b = \rho$.

*see [9].



Linear quantile regression with
Normal copula and Student/Gamma margins



ψ -linear quantile regression with
Normal copula and Student/Gamma margins

3 An open field for risk management

The bank must compute the capital needed to support the risk exposure of an operation (market, credit, operational, etc.). In general, the capital charge is determined so that the estimated probability of unexpected loss exhausting capital is less than some target insolvency rate.

3.1 General framework of capital allocation

Quantile notion of the risk Let \mathbf{F} be the (**potential**) loss probability distribution (we denote ϑ the corresponding r.v.) and $1 - \alpha$ be the target insolvency rate. The capital charge VaR (or Capital-at-Risk/Value-at-Risk) is defined by

$$\Pr \{ \vartheta > \text{VaR} \} = 1 - \alpha$$

or by

$$\text{VaR} = \inf \{ x \mid \mathbf{F}(x) \geq 1 - \alpha \}$$

In general, we distinguish Economic Capital (computed with internal models) and Regulatory Capital (computed according to methods given by the Basel Committee on Banking Supervision).

Let consider an example of equity portfolio with a long position in the security S_t . We define the loss variable as follows: $\vartheta = S_{t+1} - S_t$. We assume two distributions: $\vartheta \sim \mathcal{N}(0, 1)$ and $\vartheta \sim t_4$.

| Rating | <i>Regulatory</i> | BBB | A | AA | AAA |
|---------------------|-------------------|----------|---------|---------|----------|
| α | 99% | 99.75% | 99.9% | 99.95% | 99.97% |
| Return time | 100 days | 400 days | 4 years | 8 years | 13 years |
| $\Phi^{-1}(\alpha)$ | 2.33 | 2.81 | 3.09 | 3.29 | 3.43 |
| $t_4^{-1}(\alpha)$ | 3.75 | 5.60 | 7.17 | 8.61 | 9.83 |

Let consider now a portfolio with different securities. The Capital-at-Risk will be influenced by

- the assumption on the distributions of individual risk factors;
- and by the assumption on the **dependence** between the different risk factors.

An example (market risk) Three portfolios with five commodities of the London Metal Exchange (see [3]):

| | AL | AL-15 | CU | NI | PB |
|-------|----|-------|----|----|----|
| P_1 | 1 | 1 | 1 | 1 | 1 |
| P_2 | -1 | -1 | -1 | 1 | 1 |
| P_3 | 2 | 1 | -3 | 4 | 5 |

- Gaussian margins and Normal copula

| | 90% | 95% | 99% | 99.5% | 99.9% |
|-------|-------|-------|-------|-------|-------|
| P_1 | 7.26 | 9.33 | 13.14 | 14.55 | 17.45 |
| P_2 | 4.04 | 5.17 | 7.32 | 8.09 | 9.81 |
| P_3 | 13.90 | 17.82 | 25.14 | 27.83 | 33.43 |

- Student margins ($\nu = 4$) and Normal copula

| | 90% | 95% | 99% | 99.5% | 99.9% |
|-------|-------|-------|-------|-------|-------|
| P_1 | 6.51 | 8.82 | 14.26 | 16.94 | 24.09 |
| P_2 | 3.77 | 5.00 | 7.90 | 9.31 | 13.56 |
| P_3 | 12.76 | 17.05 | 27.51 | 32.84 | 49.15 |

3.2 Operational risk

Industry definition = “*the risk of direct or indirect loss resulting from inadequate or failed internal processes, people and systems or from external events*” (thefts, disasters, etc.).

Operational risk is now explicitly concerned by the New Basel Capital Accord (banks have to allocate capital for operational risk since 2005).

Loss Distribution Approach (LDA) Under this approach, the bank estimates, for each business line/risk type cell, the probability distributions of the severity (single event impact) and of the one year event frequency using its internal data. With these two distributions, the bank then computes the probability distribution of the aggregate operational loss. The total required capital is the sum of the *Value-at-Risk* of each business line and event type combination.

Let i and j denote a given business line and a given event type.

- $\zeta(i, j)$ is the random variable which represents the **amount of one loss event** for the business line i and the event type j . The **loss severity distribution** of $\zeta(i, j)$ is denoted by $F_{i,j}$.
- $N(i, j)$ is the random variable which represents the **number of one year events** for the business line i and the event type j . The **loss frequency distribution** of $\zeta(i, j)$ is denoted by $P_{i,j}$.

In **LDA**, the loss for the business line i and the event type j is

$$\vartheta(i, j) = \sum_{n=0}^{N(i,j)} \zeta_n(i, j)$$

The distribution $G_{i,j}$ of $\vartheta(i, j)$ is then a compound distribution

$$G_{i,j}(x) = \begin{cases} \sum_{n=1}^{\infty} p_{i,j}(n) F_{i,j}^{n*}(x) & x > 0 \\ p_{i,j}(0) & x = 0 \end{cases}$$

For a given target insolvency rate $1 - \alpha$, the capital charge corresponds to

$$\text{CaR}(i, j; \alpha) = G_{i,j}^{-1}(\alpha)$$

Computing the total capital charge The total capital charge for the bank will be then the simple summation of the capital charges accross each of the business lines and event types:

$$\text{CaR}(\alpha) = \sum_{i=1}^I \sum_{j=1}^J \text{CaR}(i, j; \alpha)$$

Problem: The Basel Committee on Banking Supervision assumes implicitly that the different losses are perfectly dependent.

Let ϑ_1 and ϑ_2 be two losses with distributions G_1 and G_2 . We denote ϑ the total loss with distribution G . We have

$$\begin{aligned}\text{CaR}(\alpha) &= G^{-1}(\alpha) \\ &= \text{CaR}_1(\alpha) + \text{CaR}_2(\alpha) \\ &= G_1^{-1}(\alpha) + G_2^{-1}(\alpha)\end{aligned}$$

It is equivalent to assume that $C\langle\vartheta_1, \vartheta_2\rangle = C^+$. In this case, we have $\vartheta_2 = G_2^{(-1)}(G_1(\vartheta_1))$. Let us denote ϖ the function $x \mapsto x + G_2^{(-1)}(G_1(x))$. We have

$$\begin{aligned}\alpha &= \Pr\{\vartheta_1 + \vartheta_2 \leq \text{CaR}(\alpha)\} \\ &= \mathbb{E}\left[\mathbf{1}_{[\varpi(\vartheta_1) \leq \text{CaR}(\alpha)]}\right] \\ &= G_1\left(\varpi^{-1}(\text{CaR}(\alpha))\right)\end{aligned}$$

It comes that $\text{CaR}(\alpha) = \varpi\left(G_1^{(-1)}(\alpha)\right)$ and we obtain the relationship

$$\text{CaR}(\alpha) = G_1^{(-1)}(\alpha) + G_2^{(-1)}\left(G_1\left(G_1^{(-1)}(\alpha)\right)\right) = \text{CaR}_1(\alpha) + \text{CaR}_2(\alpha)$$

Correlated aggregate loss distributions or correlated frequencies?

The total loss distribution ϑ for the bank as whole is defined by

$$\vartheta = \sum_{i=1}^I \sum_{j=1}^J \vartheta(i, j)$$


In this case, we could introduce the dependence directly between the aggregate loss distributions.

Or, we could introduce the dependence indirectly between the frequency distributions:


$$\vartheta = \sum_{i=1}^I \sum_{j=1}^J \sum_{n=0}^{N(i,j)} \zeta_n(i, j)$$

For example, we could use multivariate Poisson distributions generated from the Normal copula (Song [2000]).


$\xi_1 \sim \text{LN}(1,1)$ $N_1 \sim P(10)$




$\xi_2 \sim \text{LN}(1.25,0.5)$ $N_2 \sim P(12)$



Total capital charge with the Normal copula



Impact of the copula on the total loss distribution



Impact of the parameter ρ on the Capital-at-Risk

3.3 The choice of the copula function

Let consider two markets (for example the equity market and the bond market). We are interested in the probability that the loss in one market is greater than its value-at-risk given that the loss in the second market is already greater than its value-at-risk:

$$\begin{aligned}\lambda(\alpha) &= \Pr\{\vartheta_2 > \text{VaR}_2(\alpha) \mid \vartheta_1 > \text{VaR}_1(\alpha)\} \\ &= \Pr\{\vartheta_2 > \mathbf{F}_2^{-1}(\alpha) \mid \vartheta_1 > \mathbf{F}_1^{-1}(\alpha)\} \\ &= \frac{\Pr\{\mathbf{F}_2(\vartheta_2) > \alpha, \mathbf{F}_1(\vartheta_1) > \alpha\}}{\Pr\{\mathbf{F}_1(\vartheta_1) > \alpha\}} \\ &= \frac{1 - 2\alpha + \mathbf{C}(\alpha, \alpha)}{1 - \alpha}\end{aligned}$$


$\lambda(\alpha)$ depends on the copula, but not on the margins.

The limit case $\lambda = \lim_{\alpha \rightarrow 1} \lambda(\alpha)$ is called the tail dependence coefficient.


Remark 2 *The measure λ is the probability that one variable is extreme given that the other is extreme.*

1. Normal copula \Rightarrow extremes are asymptotically independent for $\rho \neq 1$, i.e $\lambda = 0$ for $\rho < 1$.
2. Student copula \Rightarrow extremes are asymptotically dependent for $\rho \neq -1$.

The copula function has then a great influence of the aggregation of risks (in particular, stress-testing is very sensitive to the choice of the copula — see [2]).



Quantile-dependent measure $\lambda(\alpha)$ for the Normal copula



$\lambda(\alpha)$ for the Student copula and $\nu = 1$

4 New pricing methods with copulas

Let consider an European Call option. The payoff is $G(T) = (S(T) - K)^+$. Under some conditions, the price $P(t_0)$ of this contingent claim is given by

$$P(t_0) = e^{-r(T-t_0)} \mathbb{E}^{\mathbb{Q}} [G(T) | \mathcal{F}_{t_0}]$$

with \mathbb{Q} the *martingale* probability measure.

For a spread option, we have $G(T) = (S_2(T) - S_1(T) - K)^+$ and we obtain a similar expression for the price.

Spread option is a special case of two-asset options. Since some years, multi-asset options are traded very frequently. The main difference with option with only one underlying is that the *martingale* probability measure is **multidimensional**.

4.1 Coherent valuation of multi-asset options

The Black-Scholes model In the BS model, we have

$$dS(t) = rS(t) dt + \sigma S(t) dW(t)$$

under \mathbb{Q} . The price of an European option is then a function of the volatility σ . However, when we compute the implied volatility from the option prices for different values of the strike K , it is not constant. This is the volatility smile effect.

Option models in banks Banks have then developed sophisticated models (e.g. stochastic volatility models) to take into account the smile effect.

To this day, therefore, the BS model continues to be used, out of analytical and computational convenience, for contingent claims based on different assets.

Problem: the margins of the multivariate Risk-Neutral Distribution (RND) are not the univariate RND.

⇒ In this case, we may show that there exists arbitrage opportunities inside the same bank (see [10]).

⇒ Moreover, one-asset options could be viewed as limits of multi-asset options — see e.g. the Basket option $G(T) = (\alpha_1 S_1(T) + \alpha_2 S_2(T) - K)^+$.

The copula construction Let \mathbb{Q} be the multivariate RND of the random vector $S(T) \mid \mathcal{F}_{t_0}$. In [10], we show that the margins of \mathbb{Q} are necessarily univariate RND*. Using Sklar's theorem, it comes that \mathbb{Q} admits the following canonical decomposition

$$\mathbb{Q}(S_1(T), \dots, S_N(T)) = \mathbf{C}^{\mathbb{Q}}(\mathbb{Q}_1(S_1(T)), \dots, \mathbb{Q}_N(S_N(T)))$$

$\mathbf{C}^{\mathbb{Q}}$ is called the risk-neutral copula (RNC).

*We prove this by using properties of the Girsanov theorem applied to multivariate probability measure.

Modelling dependence in finance using copulas

New pricing methods with copulas

Relationships between $C^{\mathbb{Q}}$ and $C^{\mathbb{P}}$

Let \mathbb{P} be the objective (or historical) distribution. We denote by $C^{\mathbb{P}}$ the objective copula. We can prove the following proposition:

Proposition 1 *If the drift and the diffusion of the asset prices vector $S(t)$ are of the form $\mu(t) \odot S(t)$ and $\sigma(t) \odot S(t)$ and if risk premiums are non stochastic, then the risk-neutral copula $C^{\mathbb{Q}}$ is equal to the objective copula $C^{\mathbb{P}}$.*

Implication of this proposition: in this case, the univariate RND can be estimated using the options market whereas the RNC can be estimated using the spot market. So, the spot market contains useful information to price multi-asset options.

The case of the spread option In [12], Valdo Durrleman shows that the price $P(t_0)$ is

$$P(t_0) = S_2(t_0) - S_1(t_0) - Ke^{-r(T-t_0)} + e^{-r(T-t_0)} \int_{-\infty}^K \int_0^{+\infty} f_1(x) \cdot \partial_1 \mathbf{C}^{\mathbb{Q}}(\mathbf{F}_1(x), \mathbf{F}_2(x+y)) \, dx \, dy$$

A remark The copula construction implies that we can associate a risk-neutral copula to a multivariate risk-neutral distribution. But it does not mean that the combination of univariate RND with a copula define necessarily a multivariate risk-neutral distribution (see [10] for further details).


BS pricing in stochastic volatility environment We assume that the asset prices $S_n(t)$ are given by the Heston model

$$\begin{cases} dS_n(t) = \mu_n S_n(t) dt + \sqrt{V_n(t)} S_n(t) dW_n^1(t) \\ dV_n(t) = \kappa_n (V_n(\infty) - V_n(t)) dt + \sigma_n \sqrt{V_n(t)} dW_n^2(t) \end{cases}$$

with $\mathbb{E} [W_n^1(t) W_n^2(t) | \mathcal{F}_{t_0}] = \rho_n (t - t_0)$, $\kappa_n > 0$, $V_n(\infty) > 0$ and $\sigma_n > 0$. The market prices of risk processes are $\lambda_n^1(t) = (\mu_n - r) / \sqrt{V_n(t)}$ and $\lambda_n^2(t) = \lambda_n \sigma_n^{-1} \sqrt{V_n(t)}$.


To compute prices of spread options, we consider that the RNC is the Normal copula with parameter ρ . We compare then the Heston prices with these given by the BS model using ATM implied volatilities.

Numerical values (two assets with same characteristics except ρ_n):
 $S_n(t_0) = 100$, $\tau = 1/12$, $r = 5\%$, $V_n(t_0) = V_n(\infty) = \sqrt{20\%}$, $\kappa_n = 0.5$,
 $\sigma_n = 90\%$ and $\lambda_n = 0$.




Volatility smile of the Heston model (one-month maturity)


$\rho = -0.5$




$\rho = 0$



$\rho = 0.5$




$\rho = 0.75$




Density of the RND of the spread ($\rho_1 = -0.75$, $\rho_2 = -0.50$)


$\rho = -0.5$




$\rho = 0$



$\rho = 0.5$




$\rho = 0.75$




Density of the RND of the spread ($\rho_1 = -0.75$, $\rho_2 = 0.50$)


Heston premium




Implied BS premium



Lower Frechet bound




Upper Frechet bound




European Spread Put premium ($\rho_1 = -0.75$, $\rho_2 = -0.50$)


Heston premium




Implied BS premium



Lower Frechet bound



Upper Frechet bound



European Spread Put premium ($\rho_1 = -0.75$, $\rho_2 = 0.50$)

How to build ‘forward-looking’ indicators for the dependence function?

In [1], Aris Bikos suggests the following method:

1. estimate the univariate RND \hat{Q}_n using Vanilla options;
2. estimate the copula \hat{C} using multi-asset options by imposing that $Q_n = \hat{Q}_n$;
3. derive “forward-looking” indicators directly from \hat{C} .

An example of the computation of the implied parameter $\hat{\rho}$

- BS model: LN distribution calibrated with ATM options; Pricing kernel = LN distributions + Normal copula


$$\hat{\rho}_1 = -0.341$$

- Bahra model: mixture of LN distributions calibrated with eight European prices; Pricing kernel = MLN distributions + Normal copula



$$\hat{\rho}_2 = 0.767$$

Remark 3 $\hat{\rho}_1$ and $\hat{\rho}_2$ are parameters of the Normal Copula. $\hat{\rho}_1$ is a Pearson correlation, not $\hat{\rho}_2$.


\Rightarrow BS model: negative dependence / Bahra model: positive dependence.



Bivariate BS density



Bivariate Bahra density



A spread option example

4.2 The pricing of credit derivatives

In multi-asset options, the risk is a market risk (because of the **volatility** of the asset prices). In credit derivatives, the risk is a credit risk (because of the **default** of the counterparties).

A default is generally described by a *survival* function $S(t) = \Pr\{T > t\}$. Let \check{C} be a *survival* copula. A multivariate survival function S can be defined as follows

$$S(t_1, \dots, t_N) = \check{C}(S_1(t_1), \dots, S_N(t_N))$$

where (S_1, \dots, S_N) are the marginal survival functions. Nelsen [1999] notices that “ \check{C} couples the joint survival function to its univariate margins in a manner completely analogous to the way in which a copula connects the joint distribution function to its margins”.

⇒ Introducing dependence between defaultable securities can then be done using the copula framework (see [21] and [25]).

Some examples*

The Default Digital Put (DDP) option

The European DDP “pays off 1 at t iff there has been a default at some time before (or including) t ”.

If we assume that the interest rate and the first default $\tau = \bigwedge_{n=1}^N T_n$ are independent, the price at time t_0 of the DDP of maturity t is then[†]

$$\begin{aligned} \text{DDP}(t_0, t) &= \mathbb{E} \left[e^{-\int_{t_0}^t r(s) ds} \mathbf{1}_{[\tau < t]} \right] \\ &= (1 - S_\tau(t)) P(t_0, t) \\ &= \left(1 - \check{\mathbf{C}}(\mathbf{S}_1(t), \dots, \mathbf{S}_N(t)) \right) P(t_0, t) \end{aligned}$$

*These examples are taken from [18].

[†] $P(t_0, t)$ is the default-free bond price.

Modelling dependence in finance using copulas

New pricing methods with copulas

The First-to-Default (FtoD) option – A first-to-default is a contingent claim that pays at the first of N credit events an amount $\varpi(\tau)$.

$$\begin{aligned} \text{FtoD}(t_0, t) &= \mathbb{E} \left[\varpi(\tau) e^{-r(\tau)} \mathbf{1}_{[\tau < t]} \right] \\ &= \int_{t_0}^t \varpi(s) e^{-r(s)} f_{\tau}(s) \, ds \\ &= \sum_{n=1}^N \int_{t_0}^t \varpi(s) e^{-r(s)} f_n(s) \partial_n \check{\mathbf{C}}(\mathbf{S}_1(s), \dots, \mathbf{S}_N(s)) \, ds \end{aligned}$$

The n^{th} -to-Default ($P_{n:N}$) option


$$\begin{aligned} P_{n:N}(t_0, t) &= \mathbb{E} \left[\varpi(T_{n:N}) e^{-r(T_{n:N})} \mathbf{1}_{[T_{n:N} < t]} \right] \\ &= \int_{t_0}^t \varpi(s) e^{-r(s)} f_{n:N}(s) \, ds \end{aligned}$$

We have of course $P_{1:N} = \text{FtoD}$ and $P_{N:N} = \text{LtoD}$ (*last-to-default*).


Numerical illustrations

$\varpi = 1$ — $r = 5\%$ — Exponential survival times with hazard rates λ_n
— the survival copula is a Normal copula with a matrix of parameters of the form


$$\begin{bmatrix} 1 & \rho & \cdots & \rho \\ & 1 & \ddots & \vdots \\ & & 1 & \rho \\ & & & 1 \end{bmatrix}$$



Influence of the parameter ρ on the DDP premium (N = 2)




Value of the FtoD premium for $\rho = 0.5$




Value of the FtoD premium for $t = 1$


N = 2




N = 3



N = 4



N = 5



Value of the premium $P_{n:N}(t_0, t)$ for $t = 1$

5 Conclusion

The use of copulas in finance is very recent.

However, private communications with professionals of other banks indicate that copulas are **largely** studied (and used) in most banks.

And professionals expect a lot from copulas to solve (and understand) many financial problems.

6 References (Copulas and Finance)

- [1] Bikos, A. [2000], Bivariate FX PDFs: a Sterling ERI application, Bank of England, *Working Paper*
- [2] Bouyé, E., V. Durrleman, A. Nikeghbali, G. Riboulet and T. Roncalli [2000], Copulas for finance — a reading guide and some applications, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [3] Bouyé, E., V. Durrleman, A. Nikeghbali, G. Riboulet and T. Roncalli [2000], Copulas: an open field for risk management, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [4] Bouyé, E., N. Gaussel and M. Salmon [2000], Investigating dynamic dependence using copulae, Financial Econometric Research Centre, City University Business School, *Working Paper*
- [5] Ceske, R. and J.V. Hernández [1999], Where theory meets practice, *Risk Magazine (Operational Risk Report)*, **12**, November, 17-20
- [6] Ceske, R., J.V. Hernández and L.M. Sánchez [2000], Quantifying event risk: the next convergence, *The Journal of Risk Finance*, **1(3)**, 9-23
- [7] Cherubini, U. and E. Luciano [2000], Multivariate option pricing with copulas, University of Turin, *Working Paper*
- [8] Cherubini, U. and E. Luciano [2000], Value at risk trade-off and capital allocation with copulas, University of Turin, *Working Paper*

- [9] Costinot, A., T. Roncalli and J. Teiletche [2000], Revisiting the dependence between financial markets with copulas, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [10] Coutant, S., V. Durrleman, G. Rapuch and T. Roncalli [2001], Copulas, multivariate risk-neutral distributions and implied dependence functions, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [11] Coutant, S., P. Martineu, J. Messines, G. Riboulet and T. Roncalli [2001], Credit risk modelling with copulas, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [12] Durrleman, V. [2001], Implied correlation and spread options, Princeton University, *Working Paper*
- [13] Durrleman, V., A. Nikeghbali and T. Roncalli [2000], How to get bounds for distribution convolutions? A simulation study and an application to risk management, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [14] Embrechts, P., A.J. McNeil and D. Straumann [1999], Correlation and dependency in risk management: properties and pitfalls, ETH Zürich, *Working Paper*
- [15] Embrechts, P., A. Hoeing and A. Juri [2001], Using copulae to bound the value-at-risk for functions of dependent risk, ETH Zürich, *Working Paper*
- [16] Frachot, A., P. Georges and T. Roncalli [2001], Loss Distribution Approach for operational risk, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [17] Frey, A. and A.J. McNeil [2000], Modelling dependent defaults, ETH Zürich, *Working Paper*

- [18] Georges, P., A-G. Lamy, E. Nicolas, G. Quibel and T. Roncalli [2001], Multivariate survival modelling: a unified approach with copulas, Groupe de Recherche Opérationnelle, Crédit Lyonnais, *Working Paper*
- [19] Giesecke, K. [2001], Structural modeling of correlated defaults with incomplete information, Humboldt-Universität zu Berlin, *Working Paper*
- [20] Hamilton, D., J. James and N. Webber [2001], Copula methods and the analysis of credit risk, University of Warwick, *Working Paper* (original version December 2000)
- [21] Li, D.X. [2000], On default correlation: a copula function approach, *Journal of Fixed Income*, **9(4)**, 43-54
- [22] Lindskog, F. [2000], Modelling dependence with copulas, ETH Zürich, *Master Thesis*
- [23] Lindskog, F. and A.J. McNeil [2001], Common poisson shock models, RiskLab, *Research Paper*
- [24] Luciano, E. and M. Marena [2001], Value at risk bounds for portfolios of non-normal returns, University of Turin, *Working Paper*
- [25] Maccarinelli, M. and V. Maggiolini [2000], The evolving practice of credit risk management in global financial institutions, *Risk Conference*, 26/27 September, Paris
- [26] Malevergne, Y. and D. Sornette [2001], General framework for a portfolio theory with non-Gaussian risks and non-linear correlations, University of Nice, *Working Paper*
- [27] Patton, A. [2000], Modelling time-varying exchange rate dependence using the conditional copula, University of California, San Diego, *Working Paper*

- [28] Rockinger, M. and E. Jondeau [2001], Conditional dependency of financial series: an application of copulas, Groupe HEC, *Cahier de Recherche*, **723/2001**
- [29] Rosenberg, J.V. [1999], Semiparametric pricing of multivariate contingent claims, Stern School of Business, *Working Paper*, **S-99-35**
- [30] Rosenberg, J.V. [2000], Nonparametric pricing of multivariate contingent claims, Stern School of Business, *Working Paper*, **FIN-00-001**
- [31] Scaillet, O. [2000], Nonparametric estimation of copulas for time series, Université Catholique de Louvain, *Working Paper*
- [32] Schönbucher, P.J. and D. Schubert [2001], Copula-dependent default risk in intensity models, Bonn University, *Working Paper*
- [33] Tibiletti L. [2000], Incremental value at risk and VaR with background-risk: traps and misinterpretations, University of Turin, *Working Paper*

7 Other references

- [1] Deheuvels, P. [1978], Caractérisation complète des lois extrêmes multivariées et de la convergence des types extrêmes, *Publications de l'Institut de Statistique de l'Université de Paris*, **23**, 1-36
- [2] Genest, C., K. Ghoudi and L-P Rivest [1995], A semiparametric estimation procedure for dependence parameters in multivariate families of distributions, *Biometrika*, **82(3)**, 543-552
- [3] Genest, C. and J. MacKay [1986], Copules archimédiennes et familles de lois bidimensionnelles dont les marges sont données, *Canadian Journal of Statistics*, **14(2)**, 145-159
- [4] Joe, H. and J.J. Xu [1996], The estimation method of inference functions for margins for multivariate models, Department of Statistics, University of British Columbia, *Technical Report*, **166**
- [5] Nelsen, R.B. [1998], An Introduction to Copulas, *Lectures Notes in Statistics*, **139**, Springer Verlag, New York
- [6] Schweizer, B. and E. Wolff [1981], On nonparametric measures of dependence for random variables, *Annals of Statistics*, **9**, 879-885
- [7] Song, P. X-K. [2000], Multivariate dispersion models generated from Gaussian copula, *Scandinavian Journal of Statistics*, **27-2**, 305-320